


Key facts

	Females	Males	Total
Number of new cases per year	252	266	518
Incidence rate (cases per 100,000 per year)	9.1	11.5	-
Cumulative lifetime risk of diagnosis (to age 74)	1 in 149	1 in 111	-
Percentage of all invasive cancers *	2.6%	2.4%	2.5%
Ranking among most common cancers *	9 th	11 th	7 th
Number of deaths per year	237	246	483
Mortality rate (deaths per 100,000 per year)	8.5	10.9	-
Cumulative lifetime risk of death (to age 74)	1 in 167	1 in 125	-
Percentage of all cancer deaths	5.8%	5.4%	5.6%
Ranking among most common invasive cancer deaths	5 th	4 th	3 rd
Number of people with this cancer still alive, end 2014 #	331	328	659
Number alive per 100,000	14	14	-

*invasive cancers only, excluding non-melanoma skin cancer


diagnosed since 1994

Cases mapped by county



Annual average number of cases, 2005-2014:
colours indicate higher or lower than
expected based on national rates


Age profile at diagnosis and death



Top risk factors


- Tobacco smoking
- Smokeless tobacco
 - Alcohol
- Chronic pancreatitis
 - Diabetes
- Greater body fatness/abdominal fatness
 - Height/tallness
- Occupational exposure to styrene
- Family history of pancreatic cancer

Trend in incidence and mortality



APC[#] Incidence: Males (94-14) 0.6%; Females (94-14) 0.5%
Mortality: Males (95-05) -1.7% (05-08) 6.4%, (08-13) -3.4%; Females (94-13) 0.3%
Annual Percentage Change

Stage at diagnosis



Treatment within 1 year

73%
pathologically
diagnosed

9%
Radiotherapy

15% Surgery

35%
Chemotherapy

% of all cases diagnosed 2012-2014

5 year net survival

Diagnosed	Net survival (age standardised)	95% Confidence interval
1994-1998	5.6%	4.3-6.8%
1999-2003	6.0%	4.8-7.2%
2004-2008	7.0%	5.8-8.2%
2009-2013	8.2%	6.4-9.9%

*10 year net survival (hybrid analysis):

7.5% (5.8-9.2%)