


Key facts

| | Females | Males | Total |
|--|---------|-----------------|-------|
| Number of new cases per year | - | 3364 | - |
| Incidence rate (cases per 100,000 per year) | - | 148.3 | - |
| Cumulative lifetime risk of diagnosis (to age 74) | - | 1 in 7 | - |
| Percentage of all invasive cancers * | - | 30.3% | - |
| Ranking among most common cancers * | - | 1 st | - |
| Number of deaths per year | - | 527 | - |
| Mortality rate (deaths per 100,000 per year) | - | 23.4 | - |
| Cumulative lifetime risk of death (to age 74) | - | 1 in 100 | - |
| Percentage of all cancer deaths | - | 11.5% | - |
| Ranking among most common invasive cancer deaths | - | 3 rd | - |
| Number of people with this cancer still alive, end 2014 # | - | 30642 | - |
| Number alive per 100,000 | - | 1344 | - |


*invasive cases only, excluding non-melanoma skin cancer

diagnosed since 1994

Cases mapped by county



Age profile at diagnosis and death




Top risk factors

- Family history of prostate cancer
 - Height/tallness
- Arsenic and arsenic compounds
- Cadmium and cadmium compounds

Annual average number of cases, 2005-2014:
 colours indicate higher or lower than
 expected based on national rates

Trend in incidence and mortality




APC[#] Incidence: (94-04) 7.5%, (04-11) 2.2%, (11-14) -3.9%


Mortality: (94-04) -0.8%, (04-13) -3.6%

Annual Percentage Change

Stage at diagnosis



Treatment within 1 year



5 year net survival

| Diagnosed | Net survival (age standardised) | 95% confidence interval |
|-----------|---------------------------------|-------------------------|
| 1994-1998 | 65.8% | 63.6-67.9% |
| 1999-2003 | 82.3% | 81.1-83.6% |
| 2004-2008 | 91.0% | 90.0-92.0% |
| 2009-2013 | 91.5% | 90.4-92.7% |

*10 year net survival (hybrid analysis):

88.7% (86.8-90.6%)