


## Key facts

	Female	Male	Total
<b>Number of new cases per year</b>	204	361	565
<b>Incidence rate (cases per 100,000 per year)</b>	7.5	15.5	-
<b>Cumulative lifetime risk of diagnosis (to age 74)</b>	1 in 179	1 in 84	-
<b>Percentage of all invasive cancers *</b>	2.1%	3.3%	2.7%
<b>Ranking among most common cancers *</b>	12 <sup>th</sup>	7 <sup>th</sup>	6 <sup>th</sup>
<b>Number of deaths per year</b>	119	193	313
<b>Mortality rate (deaths per 100,000 per year)</b>	4.2	8.6	-
<b>Cumulative lifetime risk of death (to age 74)</b>	1 in 333	1 in 167	-
<b>Percentage of all cancer deaths</b>	2.9%	4.2%	3.6%
<b>Ranking among most common invasive cancer deaths</b>	8 <sup>th</sup>	6 <sup>th</sup>	5 <sup>th</sup>
<b>Number of people with this cancer still alive, end 2014 #</b>	705	1150	1855
<b>Number alive per 100,000</b>	30	50	40


\*invasive cases only, excluding non-melanoma skin cancer  
# diagnosed since 1994

## Cases mapped by county



Annual average number of cases, 2005-2014:  
colours indicate higher or lower than  
expected based on national rates


## Age profile at diagnosis and death




## Top risk factors

- *Helicobacter pylori* infection
  - Epstein-Barr virus
  - Tobacco smoking
- Salt, salted and salty foods, or salt preserved foods, including Chinese-style salted fish
  - Ionizing radiation
- Low socio-economic status

## Trend in incidence and mortality




## Stage at diagnosis



**APC# Incidence: Males (94-03) -2.8%, (03-14) -0.2%; Females (94-14) -1.2%**  
**Mortality: Males (94-13) -3.4%; Females (94-13) -3.4%**

# Annual Percentage Change

## Treatment within 1 year



## 5 year net survival

Diagnosed	Net survival (age standardised)	95% confidence interval
1994-1998	17.4%	15.6-19.1%
1999-2003	17.3%	15.6-19.0%
2004-2008	23.2%	21.3-25.0%
2009-2013	27.4%	25.1-29.8%

\*10 year net analysis (hybrid analysis):

24.5% (21.9-27.0%)